

The number and coloration of white flank spots predict the strength of a cutaneous immune response in female Diamond Firetails, *Stagonopleura guttata*

Valeria Zanollo · Matteo Griggio · Jeremy Robertson · Sonia Kleindorfer

Received: 20 January 2012 / Revised: 11 April 2012 / Accepted: 23 April 2012 / Published online: 12 May 2012
© Dt. Ornithologen-Gesellschaft e.V. 2012

Abstract Colour variation in birds is often used to signal functional differences between individuals and sexes, though white plumage has usually been disregarded because white feathers were thought to be cheap to produce and hence unreliable signals. Here, we provide evidence for sex-specific morphological and colour differences in the strikingly patterned but apparently monomorphic Diamond Firetail (*Stagonopleura guttata*). We found that males had longer and darker tails, wider lores and a darker bill, while females had significantly more white flank spots. These gender-specific trait differences could be used to signal individual quality in sexual or social interactions, and hence we examined the correlation between these traits and the cell-mediated immune response (phytohaemagglutinin, PHA test). During both the breeding and non-breeding seasons, we found a positive correlation between PHA response and white flank spot number as well as their ultra-violet reflectance in females, and a negative correlation with white chroma and hue. In males, there was a positive correlation between PHA response and red rump reflectance (hue, red-chroma). There was no association between PHA response and either tail length or lore depth in either sex. These results add to a growing body of

evidence that female spot number and white plumage reflectance signal quality.

Keywords Sexual dimorphism · Honest signals · White plumage · Spottiness · Female quality · PHA test

Zusammenfassung

Anzahl und Farbton der weißen Seitenflecken als Indikator der kutikulären Immunantwort von Diamantfinken (*Stagonopleura guttata*)

In vielen Vogelarten dienen Farbmerkmale dazu funktionale Unterschiede zwischen Individuen und Geschlechtern zu signalisieren. Weiße Gefiederfarben sind in diesem Kontext selten untersucht worden, da angenommen wird, dass die Produktion weißer Federn mit geringen Kosten verbunden ist und somit ein unzuverlässiges Signal darstellt. In der vorliegenden Studie beschreiben wir geschlechtsspezifische Unterschiede in Morphologie und Färbung des auffallend bunten, sexuell monomorphen Diamantfinken (*Stagonopleura guttata*). Männchen besaßen längere und dunklere Schanzfedern, breitere dunkle Streifen zwischen Auge und Schnabel und einen dunkleren Schnabel, während Weibchen eine signifikant größere Anzahl weißer Seitenflecken aufwiesen. Da diese Merkmale in sexuellen oder sozialen Interaktionen genutzt werden können untersuchten wir den Zusammenhang zwischen ihnen und der zellbasierten Immunantwort (Phytohaemagglutinin, PHA test). Sowohl innerhalb als auch außerhalb der Brutsaison fanden wir eine positive Korrelation zwischen der PHA-Antwort und der Anzahl der weißen Flecken sowie der UV-Färbung der Weibchen. Die Korrelation mit Farbsättigung und Farbton der weißen Flecken war signifikant negativ. Bei den Männchen zeigte

Communicated by L. Fusani.

V. Zanollo · J. Robertson · S. Kleindorfer
School of Biological Sciences, Flinders University of South
Australia, Adelaide, SA, Australia

M. Griggio (✉)
Department of Integrative Biology and Evolution,
Konrad Lorenz Institute of Ethology,
University of Veterinary Medicine Vienna,
Savoyenstraße 1a, 1160 Vienna, Austria
e-mail: matteo.griggio@vetmeduni.ac.at

sich eine positive Korrelation zwischen PHA-Antwort und roter Färbung (Farbsättigung und Farbton) des Bürzels. In beiden Geschlechtern fanden wir keinen Zusammenhang zwischen der Länge der Schwanzfedern oder der Breite des Kopfstreifens und der PHA-Antwort. Diese Ergebnisse erweitern unser Wissen über den Gebrauch von Fleckenmuster und weißer Gefiederfarbe als weibliches Qualitätsmerkmal.

Introduction

Colour variation is a widespread phenomenon in birds and its evolution and adaptive function has been hotly debated since Darwin (reviewed in Cronin 1991). Colour variation is often correlated with sex and age (Butcher and Rohwer 1989), and hence birds may use visual cues to recognise the sex and maturity of conspecifics. Colour may also signal functional differences in quality that can be used to discriminate between potential partners and rivals, and hence optimise mate choice (McGraw and Ardia 2003; Bortolotti et al. 2006; Senar 2006, Amundsen and Pärn 2006; Clutton-Brock 2007). Curiously, despite the apparent advantages of having colour signals that can be used to discriminate between the sexes, over 50 % of bird species are monomorphic in appearance (Griffiths et al. 1998; Heinsohn et al. 2005). Yet this apparent similarity in the sexes may be deceptive because birds have a broader visual spectrum and can detect cues invisible to humans (Cuthill et al. 2000; Eaton 2005). Hence, species of birds that appear monomorphic to humans may well be dimorphic to their conspecifics, and use subtle variations in their plumage as both species-specific signals and for communication within and between the sexes.

Colour variation may also be a reliable indicator of the general health of an individual (Jawor and Breitwisch 2003; Hill and Montgomerie 1994), whereby brighter plumage and integumentary colour (bill, legs, skin and comb) have been correlated with better nutritional condition (Doucet 2002; Griffith and Pryke 2006), fewer parasites (Zuk et al. 1990; McGraw and Hill 2000; Mougeot et al. 2005) or a stronger immune response (Linville and Breitwisch 1997; Saks et al. 2003; McGraw et al. 2006; Jacquin et al. 2011). Most research on feather colouration has concentrated on melanin-based ornaments (Evans et al. 2000; Griggio and Hoi 2010; for a review, see McGraw 2006), carotenoid-based ornaments (Lonzano 1994; Hill 2002; Griggio et al. 2009a; for a review, see Hill 2006), and structural colouration (Doucet 2002; Siefferman and Hill 2005; Griggio et al. 2009b, 2010). In contrast, white plumage patches have been little studied even though they are present in many birds with different grades of

expression and conspicuousness. In part, white patches have been little researched because the keratin of unpigmented feathers is structurally achromatic and such ornaments were considered cheap to produce (Török et al. 2003; Prum 2006). However, there is growing evidence that white patches have a cost and therefore a potential function as an honest signal (Höglund and Lundberg 1987; Pärt and Qvarnström 1997; Kose and Møller 1999; Kilpimaa et al. 2004; McGlothlin et al. 2007; Griggio et al. 2011). For example, white feathers are costly in Barn Swallows (*Hirundo rustica*) because feather lice are more frequent on feathers without melanin and these unpigmented feathers are more likely to break (Kose and Møller 1999). Similarly, experimentally manipulating the diet of Dark-eyed Juncos (*Junco hyemalis*) has shown that they develop larger and brighter white patches when they have a high-protein diet (McGlothlin et al. 2007). Hence, in at least some species, it is clear that white plumage patches are costly to produce and maintain and could function as reliable signals of phenotypic quality.

Moreover, the majority of studies investigating the cost and possible function of plumage ornaments have largely focused on male ornaments, and there has been little research into females, especially in species in which the sexes are apparently equally ornamented (reviewed in Clutton-Brock 2009). In addition, there are only a few recent studies investigating the role of a white plumage patches as indicators of female quality (Velando et al. 2001; Potti and Merino 1996; Morales et al. 2007; Hanssen et al. 2009).

Our study species is the Diamond Firetail (*Stagonopleura guttata*), an endemic Australian estrildid finch that is notoriously difficult to sex in the hand (Higgins et al. 2006). Both sexes have a conspicuous plumage characterised by a red bill, black lores and breast band. The species derives its name from the conspicuous red rump and a striking pattern of white spots (diamonds) on black flanks. The carotenoid-based hue of the bill is a responsive and reliable short-term indicator of carotenoid concentration that supplements the longer-term signalling of the feathers (Stirnemann et al. 2009). We have previously shown that the number of white flank spots varies between individuals and is correlated with female (but not male) social status and dominance during feeding contests (Crowhurst et al. 2012). Exposure of the white spots also varies with behavioural context (resting, feeding, displaying) and with the intensity of behavioural interactions (Dunbar 2008; personal observation). The aim of this study is to identify which sexually dimorphic traits could be an expression of individual condition. First, we identify morphological or colour traits that are dimorphic; and second, we compare individual variation in these traits with immune activity, measured as the skin swelling to a phytohaemagglutinin

(PHA) test; this skin reaction is thought to provide information about the cell-mediated immune response (Tella et al. 2008). For the PHA response, we focus our analyses in particular on the role of variation in white spot number for response intensity.

Methods

Study species

The Diamond Firetail is a small (~17 g) estrildid finch that occurs in woodlands and open forest of temperate, semi-arid and arid areas of south-eastern Australia (Higgins et al. 2006). It is primarily granivorous and may consume the seeds of introduced grasses that have replaced native plants in much of its range (Read 1994). Since the 1980s, habitat loss has drastically reduced Diamond Firetail populations, and it is now nationally listed as near threatened (Garnett and Crowley 2000; Ford et al. 2001; Garnett et al. 2011).

Few field studies are available, and little is known about Diamond Firetail breeding or morphological variation in the wild (O’Gorman 1980; Higgins et al. 2006; McGuire and Kleindorfer 2007). Most of what is known about Diamond Firetails is based on observations by aviculturists (O’Gorman 1980, 1981; Higgins et al. 2006). The finches are socially monogamous and breed in pairs from August to February (O’Gorman 1981). The male courts the female with an elaborate display that usually involves bobbing on a branch while holding a piece of green grass. Both adults usually build the bottle-shaped nest, and females lay 3–7, but usually 4–5, eggs (Higgins et al. 2006). Both parents incubate and feed nestlings and fledglings (Higgins et al. 2006). Juveniles acquire their adult plumage at ~10–14 weeks in warmer climates and ~16–20 weeks in cooler climates; males acquire their mature plumage earlier than females (O’Gorman 1980). Both sexes have red rump feathers, black lores, black breast bands and, usually, >20 white flank spots (Higgins et al. 2006). Aviculturists assess the sex of the birds using bill colouration during the breeding season when male bills are described as “almost purple” and female bills as “coral pink” (O’Gorman 1980; Higgins et al. 2006).

Morphological measurements

We measured morphological features of captive finches: 36 males and 33 females at the Adelaide Zoo in 2006, 2007, and 2009, and 42 males and 36 females in aviaries at Flinders University between 2009 and 2011. We also measured wild finches: 14 males and 17 females caught at Monarto Zoological Park (35°8’S, 139°8’E) and 10 males

and 6 females at Sandy Creek Conservation Park (34°36’S, 138°51’E) in South Australia between 2004 and 2009. Thus, the total sample size was 102 males and 92 females. We measured six morphological traits using a digital calliper (Mitutoyo, Japan) to the nearest 0.1 mm: (1) bill length (bill tip to the base of feathers); (2) flattened wing length; (3) tail length (middle tail feather); (4) tarsus length; (5) breast band width (measured at rest, with the bird in an upright position); and (6) left and right lore width. We also measured the mass (g) with an electronic scale (EEA digital Scale, T200; Diamond Systems) to the nearest 0.1 g. All measurements were made by S.K. to eliminate measurement variation between researchers. Each bird was individually marked with an Australian Bird and Bat Banding Scheme alloy identification band and a unique combination of plastic colour bands.

Bill and plumage colouration

We measured bill and feather colouration from 62 birds (31 males, 31 females) from the Flinders University captive population in October 2010. We decided to examine sex differences during the breeding season when they are most apparent, and when these differences have most commonly been observed by aviculturists. We measured colour variables from four body regions: bill (red), rump feathers (red), and left flank spots and right flank spots (white). We measured the entire colour range (300–700 nm) using an Ocean Optic USB4000 spectrometer connected to a bifurcated fibre optic probe and an Ocean Optics LS-1 tungsten halogen light source. All the colour measurements were done on the same day, in a closed room with artificial dimmed light, so that the ambient light in the room was constant through all the measurements. Measurements were calibrated using a 99 % diffuse white-standard reference tile (WS-2) subtracting the black current (electromagnetic disturbance) from the spectrum using SpectraWin software (Top Sensor Systems, Eerbeek, Netherlands). Calibration was performed before each individual. For each bird, we took three measurements for each of the four regions, removing the probe between each measurement and averaging the three scores per area. Readings were measured with a black reference pad beneath the feathers to exclude light reflectance from alternate sources; the probe, including a probe tip to reduce ambient light interference, was held at 90° and 2 mm above the focal feather. From the resulting reflectance spectra, we quantified colour using five main standard indices: (1) hue; (2) brightness; (3) chroma; (4) red-chroma; and (5) UV-chroma (Montgomery 2006). Mean brightness was calculated as the mean summed reflectance ($R_{300-700\text{ nm}}$). Hue (λ_{R50}) was calculated as the wavelength halfway between the highest reflectance point and the lowest reflectance point

$[\lambda_{R50} = (\lambda_{Rmax} - \lambda_{Rmin})/2]$. Chroma was calculated as the difference between the highest and lowest reflectance divided by the average reflectance $[(R_{max} - R_{min})/R_{average}]$. UV-chroma was calculated as the proportion of the UV reflectance on total reflectance $(R_{300-400\text{ nm}}/R_{300-700\text{ nm}})$. Red-chroma was calculated as the percentages of total light reflected in the range 575–700 nm.

Spot number

Flank spot numbers on the left and the right sides of 161 individuals (84 males, 77 females) were counted at the time of banding, both from the two captive populations (Flinders University and Adelaide Zoo) and the wild populations (Monarto Zoological Park and Sandy Creek Conservation Park). To assess the possible effect of age on spot number, we compared spot number in 22 males and 14 females that were measured in 2009 and 2010 (of these, we also have measurements of 9 males and 6 females in 2011) at Flinders University.

Molecular sexing

Birds were genetically sexed. A blood sample of 0.01 ml was collected from each bird by jugular venipuncture using a 0.5-ml syringe (29 G $\frac{1}{2}$ " , 0.33 \times 12.7 mm) (Campbell 1995). The blood sample was stored on FTA paper (Whatman International, Cambridge, UK) and the sex was identified using molecular markers (DNA Solutions Laboratory, Melbourne) (Griffiths et al. 1998).

PHA response in captivity

The phytohaemagglutinin (PHA) response was used to estimate the T cell-mediated immune response (Salvante 2006; Tella et al. 2008). This test is commonly used in captive and wild birds (Kennedy and Nager 2006; Ardia and Schat 2008). Nevertheless, we are aware that the use of the PHA test as a measure of the activation of the cellular immune system has been questioned for many years, and some studies suggest that the reaction caused by the injection could just be a non-specific inflammatory response without T cell involvement (Martin et al. 2006; Sarv and Hórák 2009; Vinkler et al. 2012). Aware of these reservations, but in line with many studies that use the PHA response and make a case for the validity of the test in avian research, we use the PHA test as a measure of immune response; we refer to the response as the PHA response rather than the immune response. The PHA test was undertaken at Flinders University on 28 captive birds (15 males, 13 females) at the onset of the breeding season (October 2010), and on a different group of 48 captive birds (25 males, 23 females) during the non-breeding

season (April 2011). Each bird was injected in the left wing web with a 0.04-ml sample of a 5-mg solution of PHA in 1 ml physiological saline solution (PBS), equivalent to 0.2 mg PHA. If there is a response to the injected PHA, the skin will swell locally. We measured this swelling with a pressure sensitive spessimeter (accurate to 0.01 mm; Mitotoyo, Kawasaki, Kanagawa, Japan) before the injection and 24 h later, using the average of three measurements in each case. The difference between the initial and final wing web thickness was used as an index of the PHA response (Smits et al. 1999). We counted spot number and measured bill and feather colouration 1 day after the PHA test during both breeding and non-breeding seasons, and used the respective spot number and colours indices in the correlation analysis. To test for an association between plumage or bill colouration and the PHA response, we entered the different colour indices (chroma, red-chroma, UV-chroma, hue and brightness) into a principal components analysis to identify the indices that best explained the variation in each sex and season separately.

Body condition and feather lice in the field

If colour traits are honest signals of individual quality, then we predict a correlation between spot number and fitness parameters such as body condition and the prevalence of ectoparasites. Such differences in individual quality are likely to be more pronounced in wild finches because, unlike the captive birds, they are not supplied with ad libitum food and water, and they are more likely to encounter ectoparasites. Between 2004 and 2006, 28 wild individuals (12 females and 16 males) were sampled at Sandy Creek Conservation Park. We took standard morphological measurements (bill length feather, flattened wing length, tail length, tarsus length, mass), manually counted the flank spot number and recorded the presence or absence of ectoparasitic feather lice.

Ethical note

All procedures followed the Guidelines for the Use of Animals in Research (Flinders University), the legal requirements of Australia, and were approved by the Animal Welfare Committee of Flinders University (permit E235). The PHA injections were given 48 h after the birds were moved from the common aviaries to single cages to limit possible stress caused by capture; it has been shown that stress hormones can lead to suppressed PHA response (Ewenson et al. 2003). The PHA test did not have any measurable adverse effects on the birds, as they fed within 30 min after treatment. Birds remained in single aviaries for 24 h after the PHA treatment to reduce possible stress from social interactions in the common aviary.

Statistics

SPSS 18.0 for Windows was used for statistical tests with summary statistics presented as means ± SE. We used MANOVA to compare morphological variables in relation to sex and to compare colour indices from bill, spots and rump feathers between males and females. We used forward stepwise multiple regression analyses to separately test the relationship between the variation in PHA response (dependent variable) and the traits showing sexual dimorphism (tail length, lore width, spot number) and the relationship between PHA response and colour variables for each season. We took three measurements per trait; we have previously shown high repeatability of both colour measurements and the PHA response per bird (Stirnemann et al. 2009).

Results

Morphological measurements

We compared all morphology variables between the sexes using MANOVA, and found significant differences ($F_{9,126} = 3.569, P < 0.001$; Wilks' Lambda = 0.745; partial eta square = 0.25). Specifically, males had longer tails and wider lores than females (Table 1).

Bill and plumage colouration

We compared all colour indices between the sexes using MANOVA, and found significant differences ($F_{13,44} = 6.114, P < 0.001$; Wilks' Lambda = 0.356; partial eta square = 0.64). Males had higher values of bill hue, bill chroma, rump hue, rump chroma, and rump red-chroma;

only the bill spectrum was bimodal, with peaks in the UV (300–400 nm) and red (575–700 nm). Flank spot feathers showed similar spectra between the sexes (Table 2). Also, spot number did not differ significantly between the right and left flank in either sex (Table 2).

Spot number

Spot number differed significantly between the sexes (Table 1). Females had more spots than males, but there was overlap between the sexes (Fig. 1). We tested for a correlation between spot number and body condition (the residuals of mass against tarsus length) and found no significant correlation in either sex (Pearson correlation: males, $r = -0.01, P = 0.957, n = 68$; females, $r = 0.23, P = 0.060, n = 67$). There was no significant difference in spot number between captive or wild populations. To test this, we compared the total spot number between the combined population of captive finches held at Adelaide Zoo and Flinders University and the combined wild populations at Monarto Zoological Park and Sandy Creek Conservation Park (ANOVA: $F_{1,159} = 0.17, P = 0.678$).

Spot number did not change significantly across years (2009 and 2010) in males (paired t test: Year 1: 56.68 ± 2.53 ; Year 2: 59.59 ± 2.034 ; $t_{21} = -1.97, P = 0.061$) or females (paired t test: Year 1: 63.21 ± 2.67 ; Year 2: 65.29 ± 1.94 ; $t_{13} = -0.75, P = 0.47$). For a subsample of 15 individuals, we could test the difference in spot number over three consecutive years (2009, 2010, and 2011) and again, found no significant difference in males ($n = 9$) or females ($n = 6$) (Friedman Test: males, $\chi^2 = 0.76, P = 0.682$; females, $\chi^2 = 3.00, P = 0.220$). There was no significant difference in the residuals of the regression between spot number in Year 1 and Year 2 for either sex ($P > 0.6$).

Table 1 Morphology and spot number in male and female Diamond Firetails (*Stagonopleura guttata*)

Trait	Male		Female		F	df	P
	Range	Mean ± SE	Range	Mean ± SE			
Bill length	8.7–13.9	11.15 ± 0.10	8.7–12.8	10.92 ± 0.09	1.02	133	0.314
Wing length	61–71	66.23 ± 1.16	61–70	65.9 ± 1.83	1.62	133	0.205
Tail length	30–43	39.61 ± 0.23	34–46	38.72 ± 0.23	5.17	133	0.025
Tarsus	15.4–19.0	17.17 ± 0.73	13.9–19.8	16.88 ± 0.09	3.50	133	0.063
Mass	14–28	18.33 ± 0.20	16–24	18.39 ± 0.21	0.003	133	0.955
Breast band	7–23	13.31 ± 0.39	6.7–20.2	12.78 ± 0.40	0.25	133	0.618
Lore width	3.75–5.90	4.82 ± 0.47	3.65–5.63	4.61 ± 0.05	9.26	133	0.003
N. Spot (left)	16–44	27.82 ± 0.74	17–44	32.36 ± 0.69	16.25	133	<0.001
N. Spot (right)	11–51	27.90 ± 0.77	18–45	32.68 ± 0.66	21.25	133	<0.001
N. Spot (total)	33–93	55.63 ± 1.40	36–83	65.04 ± 1.26	21.48	133	<0.001

Statistical results are shown for a MANOVA test (male, $n = 66$; female, $n = 69$), with P values <0.05 shown in bold

Table 2 Bill and feather colouration in male and female captive Diamond Firetails

Trait	Male		Female		<i>F</i>	<i>df</i>	<i>P</i>
	Range	Mean ± SE	Range	Mean ± SE			
Bill hue	620.46–648.52	634.50 ± 1.33	594.39–641.54	620.38 ± 2.05	34.34	56	<0.001
Bill brightness	10.05–31.47	19.77 ± 0.95	11.70–27.07	18.32 ± 0.73	0.75	56	0.391
Bill chroma	0.66–1.85	1.10 ± 0.05	0.49–1.46	0.92 ± 0.04	6.32	56	0.015
Bill UV-chroma	0.17–0.23	0.21 ± 0.03	0.19–0.24	0.21 ± 0.02	0.51	56	0.480
Bill red-chroma	0.37–0.49	0.42 ± 0.01	0.38–0.48	0.42 ± 0.01	0.004	56	0.950
Spot hue	326.45–344.77	334.37 ± 0.88	325.18–347.98	336.16 ± 1.05	1.46	56	0.231
Spot brightness	28.29–46.81	36.83 ± 0.92	27.012–43.99	34.61 ± 0.78	3.51	56	0.066
Spot chroma	0.48–0.95	0.75 ± 0.02	0.51–1.01	0.75 ± 0.07	0.09	56	0.763
Spot UV-chroma	0.16–0.22	0.19 ± 0.01	0.16–0.21	0.19 ± 0.01	0.18	56	0.671
Rump hue	606.69–618.55	611.38 ± 0.54	595.55–617.60	607.08 ± 1.01	8.89	56	0.004
Rump chroma	2.93–4.14	3.72 ± 0.05	2.35–3.94	3.44 ± 0.07	10.02	56	0.003
Rump red-chroma	0.75–0.93	0.88 ± 0.01	0.69–0.93	0.85 ± 0.01	7.36	56	0.009

Statistical results are shown for a MANOVA test (male, $n = 30$; female, $n = 28$), with P values <0.05 shown in bold




Fig. 1 Spot number on the *right* and *left* flanks of male (open circles) and female (black circles) Diamond Firetails (*Stagonopleura guttata*)

PHA response in captivity

For experimental birds used in the PHA test, there was no significant difference in spot number between the breeding and non-breeding season (ANOVA: season: $F_{1,72} = 0.003$, $P = 0.96$; sex \times season: $F_{1,72} = 3.47$, $P = 0.067$). However, as expected, spot number was significantly different between the sexes for the two experimental groups, with females having more spots (sex: $F_{1,72} = 4.97$, $P = 0.029$). The PHA response was comparable between males and females, and there was no significant effect of season (ANOVA: sex: $F_{1,72} = 0.16$, $P = 0.69$; season: $F_{1,72} = 0.03$, $P = 0.863$; sex \times season: $F_{1,72} = 0.035$, $P = 0.851$).

We compared PHA response in relation to tail length, lore width, and spot number (dimorphic traits) during the breeding and non-breeding seasons. During the breeding




Fig. 2 Relationship between PHA response and total spot number in males (open circles, solid regression line) and females (black circles, dashed regression line) during the breeding season. There was a significant relationship for females ($r = 0.69$, $n = 13$, $P = 0.009$) but not males ($r = -0.06$, $n = 15$, $P = 0.821$)

season, PHA response in females was predicted by spot number ($F_{1,12} = 10.01$, $r = 0.69$, $P = 0.009$), but not by tail length ($r = -0.30$, $P = 0.336$) or lore width ($r = 0.20$, $P = 0.539$). In males, PHA response was not related to tail length, lore width or spot number during either the breeding season or the non-breeding season (Fig. 2). During the non-breeding season, PHA response varied with spot number in females ($F_{1,22} = 6.52$, $r = 0.49$, $P = 0.018$; but not tail length: $r = -0.11$, $P = 0.617$; or lore width: $r = -0.31$, $P = 0.166$) (Fig. 3).

We used forward stepwise multiple regression to test PHA response in relation to the PCA-derived colour variables (factor loadings and cumulative variance shown in Tables 3 and 4) that were sampled from the bill, flank, spots




Fig. 3 Relationship between PHA response and total spot number in males (open circles, solid regression line) and females (black circles, dashed regression line) during non-breeding season. There was a significant relationship for females ($r = 0.49, n = 23, P = 0.018$) but not males ($r = 0.24, n = 25, P = 0.237$)

and rump. During the breeding season, male PHA response was positively correlated with PC1 scores for rump colour ($F_{1,13} = 6.60, r = 0.58, P = 0.023$) (Fig. 4) and female PHA response was negatively correlated with PC1 scores for spot colour ($F_{1,10} = 6.97, r = -0.56, P = 0.025$) (Fig. 5). That is, males with a redder rump (higher hue and red-chroma values) had a higher PHA response, and females with higher UV reflectance of the white spots and lower values of hue and chroma had a higher PHA response. During the non-breeding season, there was no effect of any colour variable on PHA response in males ($F_{4,8} = 0.33, P = 0.849$) or females ($F_{4,6} = 0.76, P = 0.584$).

Body condition and feather lice in the field

Body condition (calculated as the residual of mass against tarsus) was significantly related to total spot number

in females (Pearson correlation, $r = 0.887, P < 0.001, n = 9$), but not males (Pearson correlation, $r = 0.368, P = 0.215, n = 12$) (Fig. 6). Females with more white flank spots had higher body condition. Birds were commonly infested with feather lice; there was no significant difference in the prevalence of feather lice in females (4 infested, 8 uninfested; 33 %) and males (9 infested, 7 uninfested; 56 %) (Friedman Test: $\chi^2 = 0.317, P = 0.576$). There was also no significant relationship between feather lice infestation and the total spot number (independent t test: males, $t = -0.51, P = 0.622$; females, $t = 2.08, P = 0.071$), or body condition (independent t test: males, $t = -0.50, P = 0.631$; females, $t = 0.87, P = 0.407$).

Discussion

Our principal findings were: (1) evidence of sexual dimorphism in the apparently monomorphic Diamond Firetail; (2) in captive females, PHA response was correlated with spot number and spot colour; in captive males, PHA response was correlated with rump colour; and (3) in a wild population, female spot number was positively associated with body condition. Some of the sex differences in morphological and colour traits fit the standard expectation for species with choosy females: namely, males are more ornamented in traits that could signal quality and that could be used for female mate choice (Andersson 1994; but see Watson and Simmons 2010). In this study, males had longer tails and wider lores than females, and on average, males also had higher colour reflectance (bill, rump, flank spots) and higher R_{50} reflectance and chroma (bill, rump). We also found that bills reflect UV during the breeding season (there was no sex difference). Notably, the sex differences in bill colour reflectance and bill chroma corroborate the assessment of aviculturists in their attempts to sex Diamond Firetails by ocular estimation (O’Gorman 1980; Higgins et al. 2006).

Table 3 Unrotated factor loadings (component matrix) from principal components analysis (PCA) of the different colour indices of the bill, flank spots and rump in male ($n = 15$) and female ($n = 13$) Diamond Firetails during the breeding season

	Males						Females					
	Bill		Flank spots		Rump		Bill		Flank spots		Rump	
	PC1	PC2	PC1	PC2	PC1	PC2	PC1	PC2	PC1	PC2	PC1	PC2
	61 %	25.2 %	67.8 %	29 %	70.6 %	25.3 %	61.4 %	22.7 %	58.5 %		71 %	26.1 %
Chroma	0.94		0.90	0.39	0.99		0.96		0.89		0.98	
UV-chroma	-0.78	0.35	-0.97		n.a.	n.a.	-0.87		-0.98		n.a.	n.a.
Red-chroma	0.95		n.a.	n.a.	0.97		0.90		n.a.	n.a.	0.91	0.37
Hue		0.95	0.92		0.93		0.76		0.71		0.95	
Brightness	-0.78	0.37	-0.32	0.93		0.98		0.98				0.91

Percentage values indicate cumulative variance for each component. Loadings below 0.20 are not reported




n.a. not available

Table 4 Unrotated factor loadings (component matrix) from the principal components analysis (PCA) of the different colour indices of the bill, flank spots and rump in male ($n = 13$) and female ($n = 12$) Diamond Firetails during the non-breeding season

	Males						Females					
	Bill		Flank spots		Rump		Bill		Flank spots		Rump	
	PC1	PC2	PC1	PC2	PC1	PC2	PC1	PC2	PC1	PC2	PC1	PC2
	63.8 %		46.9 %		74.8 %		59.3 %		70.4 %		81.2 %	
Chroma	0.98		-0.97		0.95		0.90		0.93		0.99	
UV-chroma	-0.86		0.81	0.54	n.a.	n.a.	-0.91		-0.72		n.a.	n.a.
Red-chroma	0.95		n.a.	n.a.	0.81		0.86		n.a.	n.a.	0.97	
Hue				0.91	0.82		0.57	0.62	0.92		0.73	
Brightness	-0.64		0.52	-0.76	0.87		-0.50	0.71	-0.76		0.89	

Percentage values indicate cumulative variance for each component. Loadings below 0.20 are not reported

n.a. not available

**Fig. 4** Relationship between PHA response and PC1 rump reflectance scores in males during the breeding season ($r = 0.58$, $n = 15$, $P = 0.023$)**Fig. 5** Relationship between PHA response and PC1 scores (UV reflectance of white flank spots) in females during the breeding season ($r = -0.56$, $n = 13$, $P = 0.027$)**Fig. 6** Relationship between body condition and total spot number in males (*black circles, solid regression line*) and females (*open circles, dashed regression line*) in a wild population. Body condition was calculated as the residual of mass against tarsus. There was a significant relationship for females ($r = 0.89$, $n = 9$, $P < 0.001$) but not males ($r = -0.37$, $n = 13$, $P = 0.215$)

A previous experimental study—involving carotenoid depletion—showed that the bill colour of Diamond Firetails in the visible range (400–700 nm) was dependant on carotenoids, and males with brighter bills had higher PHA response than duller individuals (Stirnemann et al. 2009). We were unable to repeat this finding in either the breeding or non-breeding season, possibly because birds were not carotenoid depleted; in this study, bill colouration was not related to PHA response. Carotenoids deposited in the bill are defined as mobile carotenoids because they can be easily displaced to activate the immune system and scavenge free-radicals to protect tissues from oxidative damage (Goodwin 1986; McGraw and Ardia 2003; McGraw et al. 2006). Thus, in the 24 h following the injection of PHA, there may have been a rapid reallocation of carotenoids from the bill leading to a decrease in carotenoid-based

colouration (Faivre et al. 2003) and plasma carotenoid levels (Biard et al. 2009). Carotenoids deposited in the feathers are fixed and cannot be mobilised or rapidly changed. We suggest that plumage colouration is a more reliable signal of the immune function. Indeed, we showed that males with redder rumps, a carotenoid-based plumage (McGraw and Schuetz 2004), had a stronger PHA response during the breeding season, which concurs with the findings in other species that individuals with brighter carotenoid-based plumage usually have a higher PHA response or stronger response to parasite infection (Saks et al. 2003; Blount et al. 2003; Mougeot et al. 2009).

Contradicting the convention of more ornamented males, female Diamond Firetails had more flank spots than males. This finding concurs with previous research showing that female birds are generally more spotted than males (Roulin 2003; Swaddle and Witter 1995). Significantly, we found that spot number did not vary with age, which rejects the explanation that sex-related annual mortality explains the finding of sex differences in spot number. But there were sex differences in PHA response: females with more flank spots had a higher PHA response, whereas males did not. Perhaps spot number is a reliable cue for the individual quality of females, as found in other studies. For example, in Common Starlings (*Sturnus vulgaris*), females with more spots on their breasts during the breeding season had earlier ovarian development and thus laid clutches earlier in the season (Swaddle and Witter 1995). Similarly, female Barn Owls (*Tyto alba*) with more black spots had higher parasite resistance (Roulin et al. 2001) and their offspring had increased humoral immuno-competence (Roulin 2004). Our finding of a significant relationship between spot number and PHA response in captive Diamond Firetails is potentially confounded by the abundant food resources and lack of predators and parasites in captivity. Yet, we found that wild Diamond Firetails also showed a positive and strong correlation between female spot number and body condition, supporting the idea that spot number signals individual quality. Finally, we have previously shown, in experimental feeding trials, that female Diamond Firetails with many spots win food contests in social dominance trials, whereas male spot number was not related to feeding dominance (Crowhurst et al. 2012).

We address the colour characteristics of the white spots to gain more insight into their possible signalling function. Previous research tended to view white spots as structural achromatic plumage that is cheap to produce and maintain (Török et al. 2003; Prum 2006). But we show here that birds with white spots, that had high UV reflectance, low chroma and low hue, had a higher PHA response. This association between white plumage and individual quality has been found in few other species. Female Common Eiders (*Somateria mollissima*) with fewer white feathers

did not respond to a diphtheria toxoid immune challenge (Hanssen et al. 2006). Similarly, in the South Polar Skua (*Catharacta maccormicki*), the whiteness of the male wing patch was positively related to the response against tetanus (Hanssen et al. 2009). However, these studies only considered brightness, chroma and hue, while here we also showed that females with a higher UV reflectance during the breeding season had a higher PHA response. To our knowledge, this is the first evidence that white UV-chroma correlates with the PHA response, and therefore possibly individual quality in females.

Perhaps the expression of white spots is under genetic control and is not a condition-dependent trait. A genetic mechanism for black spot number has been shown in Barn Owls (Roulin et al. 1998, 2000). Such a genetic mechanism for the expression of white spots could have pleiotropic effects on immunocompetence and other aspects of condition. If white spot number is genetically determined, then what maintains variation in spot number in the population? Why is there not extreme directional selection for males and females with many spots? One hypothesis to explain the observed variation in spot number (between individuals and sexes) is that birds with low or high spot number, for example, pursue different behavioural strategies during foraging or social contexts, and that a mix of strategies confers higher individual fitness payoffs (Roulin 2004; Mckinnon and Pierotti 2010). Our measurements showed no difference across years in adult birds. But spot number could nonetheless be a condition-dependent trait if the variation results from stress during growth and development—which needs to be experimentally tested between the nestling to yearling phase. Finally, it is also possible that spot number is highly heritable whereas UV-reflectance is strongly condition-dependent (Badayaev et al. 2001; Johnsen et al. 2003). These ideas also require future testing.

In conclusion, we have shown that both the number and reflectance of white flank spots correlate positively with female PHA response during the breeding season. Both cues could be signals of female quality. Further research should test the “dual utility hypothesis” of white spots as armament and ornament (Berglund et al. 1996; Griggio et al. 2007). White flank spots can also be used in a multiple signal system (Johnstone 1996; Badayaev et al. 2001), where number (size) and reflectance could indicate different aspects of female quality. Variation in female white spots could function in male mate choice as well as in intrasexual feeding contests and social dominance (Andersson et al. 2002; Doucet and Montgomerie 2003; Wilson et al. 2010; Crowhurst et al. 2012). Finally, the findings add support for a growing body of evidence that white plumage ornaments can be a reliable cue of individual quality (Morales et al. 2007; Hanssen et al. 2009; Lehikoinen et al. 2010).

Acknowledgments We thank the Australian Research Council, Adelaide Zoological Society, South Australian Department for Environment and Heritage, and Conservation Council of South Australia for funding. We thank the Adelaide Zoo for allowing us to measure their captive birds. We thank David Paton for stimulating discussion and assistance with observing and banding Diamond Firetails at Monarto Zoological Park. We thank Wesley M. Hochachka and two anonymous referees for their constructive and helpful comments on a previous version of the manuscript. All procedures complied with the current laws of the country in which they were performed and followed the Guidelines for the Use of Animals in Research (Flinders University) and were approved by the Animal Welfare Committee of Flinders University (permit E190 and E281).

References

- Amundsen T, Pärn H (2006) Female coloration: review of functional and non-functional hypotheses. In: Hill GE, McGraw KJ (eds) Bird coloration, vol 2: function and evolution. Harvard University Press, Cambridge, pp 280–345
- Andersson M (1994) Sexual selection. Princeton University Press, Princeton
- Andersson S, Pryke SF, Örnborg J, Lawes MJ, Andersson M (2002) Multiple receivers, multiple ornaments, and a trade-off between agonistic and epigamic signalling in a widowbird. *Am Nat* 160:683–691
- Ardia DR, Schat KA (2008) Ecoimmunology. In: Davison F, Kaspers B, Schat KA (eds) Avian immunology. Academic, London, pp 421–441
- Badayaev AV, Hill GE, Dunn PO, Glenn JC (2001) Plumage color as a composite trait: developmental and functional integration of sexual ornamentation. *Am Nat* 158:221–235
- Berglund A, Bisazza A, Pilastro A (1996) Armaments and ornaments: an evolutionary explanation of traits of dual utility. *Biol J Linn Soc* 58:385–399
- Biard C, Hardy C, Motreuil S, Moreau J (2009) Dynamics of PHA-induced immune response and plasma carotenoids in birds: should we have a closer look? *J Exp Biol* 212:1336–1343
- Blount JD, Metcalfe NB, Birkhead TR, Surai PF (2003) Carotenoid modulation of immune function and sexual attractiveness in zebra finch. *Science* 300:125–127
- Bortolotti G, Blas J, Negro J, Tella J (2006) A complex plumage pattern as an honest social signal. *Anim Behav* 72:423–430
- Butcher GS, Rohwer S (1989) The evolution of conspicuous and distinctive coloration for communication in birds. In: Power DM (ed) Current Ornithology, vol 7, Plenum, New York, pp 51–108
- Campbell TF (1995) Avian haematology and cytology, 2nd edn. Iowa State University Press, Iowa
- Clutton-Brock T (2007) Sexual selection in males and females. *Science* 318:1882–1885
- Clutton-Brock T (2009) Sexual selection in females. *Anim Behav* 75:3–11
- Cronin H (1991) The ant and the peacock. Cambridge University Press, Cambridge
- Crowhurst CJ, Zanollo V, Griggio M, Robertson J, Kleindorfer S (2012) White flank spots signal feeding dominance in female diamond firetails, *Stagonopleura guttata*. *Ethology* 118:63–75
- Cuthill IC, Partridge JC, Bennet ATD (2000) Ultraviolet vision in birds. *Adv Stud Behav* 29:159–214
- Doucet SM (2002) Structural plumage coloration, male body size, and condition in the blue-back grassquit. *Condor* 104:30–38
- Doucet MD, Montgomerie R (2003) Multiple sexual ornaments in satin bowerbirds: ultraviolet plumage and bowers signal different aspects of male quality. *Behav Ecol* 14:503–509
- Dunbar A (2008) Flank spot displays in the diamond firetail (*Stagonopleura guttata*): evidence for signal function in social contests? Honour's thesis, Flinders University, Adelaide
- Eaton MD (2005) Human vision fails to distinguish widespread sexual dichromatism among sexually “monochromatic” birds. *Proc Natl Acad Sci USA* 102:10942–10946
- Evans MR, Goldsmith AR, Norris SRA (2000) The effects of testosterone on antibody production and plumage coloration in male house sparrows *Passer domesticus*. *Behav Ecol Sociobiol* 47:156–163
- Ewenson E, Zann R, Flannery G (2003) PHA immune response assay in captive zebra finches is modulated by activity prior to testing. *Anim Behav* 66:797–800
- Favre B, Gregoire A, Preault M, Cezilly F, Sorci G (2003) Immune activation rapidly mirrored in secondary sexual traits. *Science* 300:103
- Ford H, Barrett G, Saunders D, Recher H (2001) Why have birds in the woodlands of Southern Australia declined? *Biol Conserv* 97:71–88
- Garnett S, Crowley G (2000) The action plan for Australian birds 2000. Environment Australia, Canberra
- Garnett S, Szabo J, Dutson G (2011) The action plan for Australian birds 2010. CSIRO Publishing, Melbourne
- Goodwin TW (1986) Metabolism, nutrition, and function of carotenoids. *Annu Rev Nutr* 6:273–297
- Griffith SC, Pryke SR (2006) Benefits to females of assessing color displays. In: Hill GE, McGraw KJ (eds) Bird coloration, vol 2, function and evolution. Harvard University Press, Cambridge, pp 233–279
- Griffiths R, Double M, Orr K, Dawson R (1998) A DNA test to sex most birds. *Mol Ecol* 7:1071–1075
- Griggio M, Hoi H (2010) Only females in poor condition display a clear preference and prefer males with an average badge. *BMC Evol Biol* 10:261
- Griggio M, Serra L, Licheri D, Monti A, Pilastro A (2007) Armaments and ornaments in the rock sparrow: a possible dual utility of a carotenoid-based feather signal. *Behav Ecol Sociobiol* 61:423–433
- Griggio M, Devigili A, Hoi H, Pilastro A (2009a) Female ornamentation and directional male mate preference in the rock sparrow. *Behav Ecol* 20:1072–1078
- Griggio M, Serra L, Licheri D, Campomori C, Pilastro A (2009b) Molt speed affects structural feather ornaments in the blue tit. *J Evol Biol* 22:782–792
- Griggio M, Zanollo V, Hoi H (2010) UV plumage color is an honest signal of quality in male budgerigars. *Ecol Res* 25:77–82
- Griggio M, Valera F, Casas-Crivillè A, Hoi H, Barbosa A (2011) White tail markings as indicator of quality and affect mate preference in rock sparrow. *Behav Ecol Sociobiol* 65:655–664
- Hanssen SA, Folstad I, Erikstad KE (2006) White plumage reflects individual quality in female eiders. *Anim Behav* 71:337–343
- Hanssen SA, Bustnes JO, Tveraa T, Hasselquist D, Varpe Ø, Henden JA (2009) Individual quality and reproductive effort mirrored in white wing plumage in both sexes of south polar skuas. *Behav Ecol* 20:961–966
- Heinsohn R, Legge S, Endler JA (2005) Extreme reversed sexual dichromatism in a bird without sex role reversal. *Science* 309:617–619
- Higgins P, Peter J, Cowling S (eds) (2006) Handbook of Australian, New Zealand and Antarctic Birds. Vol 7, Boatbill to starlings, part B: Dunnock to starlings. Oxford University Press, Melbourne
- Hill GE (2002) A red bird in a brown bag: the function and evolution of colourful plumage in the house finch. Oxford University Press, New York
- Hill GE (2006) Environmental regulation of ornamental coloration in bird coloration: mechanics and measurements. In: Hill GE,

- McGraw KJ (eds) Bird coloration, vol 1, mechanisms and measurements. Harvard University Press, Cambridge, MA, pp 507–560
- Hill GE, Montgomerie R (1994) Plumage colour signals nutritional condition in the house finch. *Proc R Soc Lond B* 258:47–52
- Höglund J, Lundberg A (1987) Sexual selection in a monomorphic lek-breeding bird: correlates of male mating success in the great snipe *Gallinago media*. *Behav Ecol Sociobiol* 21:211–216
- Jacquin L, Lenouvel P, Haussy C, Ducatez S, Gasparini J (2011) Melanin-based coloration is related to parasite intensity and cellular immune response in an urban free living bird: the feral pigeon *Columba livia*. *J Avian Biol* 42:11–15
- Jawor J, Breitwisch R (2003) Melanin ornaments, honesty and sexual selection. *Auk* 120:249–265
- Johnsen A, Delhey K, Andersson S, Kempenaers B (2003) Plumage colour in nestling blue tits: sexual dichromatism, condition dependence and genetic effects. *Proc R Soc Lond B* 270:1263–1270
- Johnstone RA (1996) Multiple displays in animal communication: ‘backup signals’ and ‘multiple messages’. *Philos Trans R Soc Lond B* 351:329–338
- Kennedy MW, Nager RG (2006) The perils and prospects of using phytohaemagglutinin in evolutionary ecology. *Trends Ecol Evol* 21:653–655
- Kilpimaa J, Alatalo RV, Siitari H (2004) Trade-offs between sexual advertisement and immune function in the pied flycatcher (*Ficedula Hypoleuca*). *Proc R Soc Lond B* 271:245–250
- Kose M, Møller AP (1999) Sexual selection, feather breakage, and parasites: the importance of white spots in the tail of the barn swallow (*Hirundo rustica*). *Behav Ecol Sociobiol* 45:430–436
- Lehikoinen A, Jaatinen K, Öst M (2010) Do female ornaments indicate quality in eider ducks? *Biol Lett* 6:225–228
- Linville S, Breitwisch R (1997) Carotenoid availability and plumage coloration in a wild population of northern cardinals. *Auk* 114:796–800
- Lonzano GA (1994) Carotenoids, parasites, and sexual selection. *Oikos* 70:309–311
- Martin LB, Han P, Lewittes J, Kuhlman JR, Klasting KC, Wikelski M (2006) Phytohemagglutinin-induced skin swelling in birds: histological support for a classic immunoeological technique. *Funct Ecol* 20:290–299
- McGlothlin JW, Joel W, Duffy DL, Henry-Freeman JL, Ketterson ED (2007) Diet quality affects an attractive white plumage pattern in dark-eyed juncos (*Junco hyemalis*). *Behav Ecol Sociobiol* 61:1391–1399
- McGraw KJ (2006) Mechanics of melanin-based coloration. In: Hill GE, McGraw KJ (eds) Bird coloration. Vol 1, mechanism and measurements. Harvard University Press, Cambridge, pp 243–294
- McGraw KJ, Ardia D (2003) Carotenoids, immunocompetence, and the information content of sexual colors: an experimental test. *Am Nat* 162:704–712
- McGraw KJ, Hill GE (2000) Differential effects of endoparasitism on the expression of carotenoid- and melanin-based ornamental coloration. *Proc R Soc Lond B* 267:1525–1531
- McGraw KJ, Schuetz JG (2004) The evolution of carotenoid coloration in estrild finches: a biochemical analysis. *Comp Biochem Phys B* 139:45–51
- McGraw KJ, Crino O, Medina-Jerez W, Nolan P (2006) Effect of dietary carotenoid supplementation on food intake and immune function in songbird with no carotenoid coloration. *Ethology* 112:1209–1216
- McGuire A, Kleindorfer S (2007) Nesting success and apparent nest-adornment in diamond firetails (*Stagonopleura guttata*). *Emu* 107:44–51
- Mckinnon JS, Pierotti MER (2010) Colour polymorphism and correlated characters: genetic mechanisms and evolution. *Mol Ecol* 19:5101–5125
- Montgomerie R (2006) Cosmetic and adventitious colors. In: Hill GE, McGraw KJ (eds) Bird coloration. Vol 1, mechanisms and measurements. Harvard University Press, Cambridge, pp 399–427
- Morales J, Moreno J, Merino S, Sanz JJ, Tomas G, Arriero E, Lobato E, Martinez-De La Puente J (2007) Female ornaments in the pied flycatcher *Ficedula hypoleuca*: associations with age, health and reproductive success. *Ibis* 149:245–254
- Mougeot F, Redpath S, Leckie F (2005) Ultra-violet reflectance of male and female red grouse, *Lagopus lagopus scoticus*: sexual ornaments reflect nematode parasite intensity. *J Avian Biol* 36:203–209
- Mougeot F, Pérez-Rodríguez L, Sumozas N, Terraube J (2009) Parasites, condition, immune responsiveness and carotenoid-based ornamentation in male red-legged partridge *Alectoris rufa*. *J Avian Biol* 40:67–74
- O’Gorman B (1980) A breeding study of the diamond firetail. *Aust Avic* 34:13–19
- O’Gorman B (1981) A prolonged field study of the diamond firetail. *Aust Avic* 35:14–27
- Pärt T, Qvarnström A (1997) Badge size in Collared Flycatchers predicts outcome of male competition over territories. *Anim Behav* 54:893–899
- Potti J, Merino S (1996) Decreased levels of blood trypanosome infection correlate with female expression of a male secondary sexual trait: implications for sexual selection. *Proc R Soc Lond B* 263:1199–1204
- Prum RO (2006) Anatomy, physics and evolution of structural colors. In: Hill GE, McGraw KJ (eds) Bird coloration. Vol 1, mechanism and measurements. Harvard University Press, Cambridge, pp 295–353
- Read J (1994) The diet of three species of firetail finches in temperate South Australia. *Emu* 94:1–8
- Roulin A (2003) Geographic variation in sexually selected traits: a role for direct selection or genetic correlation? *J Avian Biol* 34:251–258
- Roulin A (2004) Proximate basis of the covariation between a melanin-based female ornament and offspring quality. *Oecologia* 140:668–675
- Roulin A, Richner H, Ducrest AL (1998) Genetic, environmental, and condition-dependent effects on female and male ornamentation in the barn owl *Tyto alba*. *Evolution* 52:1451–1460
- Roulin A, Jungi TW, Pester H, Dijkstra C (2000) Female barn owls (*Tyto alba*) advertise good genes. *Proc R Soc Lond B* 267:937–941
- Roulin A, Riols C, Dijkstra C, Ducrest A (2001) Female plumage spottiness signals parasite resistance in the barn owl (*Tyto alba*). *Behav Ecol* 12:103–110
- Saks L, Ots I, Hörak P (2003) Carotenoid-based plumage coloration of male greenfinches reflects health and immunocompetence. *Oecologia* 134:301–307
- Salvante KG (2006) Techniques for studying integrated immune function in birds. *Auk* 123(2):575–586
- Sarv T, Hörak P (2009) Phytohaemagglutinin injection has a long-lasting effect on immune cells. *J Avian Biol* 40:569–571
- Senar JC (2006) Color displays as intrasexual signals of aggression and dominance. In: Hill GE, McGraw KJ (eds) Bird coloration. Vol 2, function and evolution. Harvard University Press, Cambridge, pp 87–137
- Siefferman L, Hill GE (2005) Evidence for sexual selection on structural plumage coloration in female eastern bluebirds (*Sialia sialis*). *Evolution* 59:1819–1828

- Smits JE, Bortolotti GR, Tella JL (1999) Simplifying the phytohaemagglutinin skin-testing technique in studies of avian immunocompetence. *Funct Ecol* 13:567–572
- Stirnemann I, Johnston G, Robertson J, Kleindorfer S (2009) Phytohaemagglutinin (PHA) response and bill-hue wavelength increase with carotenoid supplementation in diamond firetails (*Stagonopleura guttata*). *Emu* 109:344–351
- Swaddle J, Witter M (1995) Chest plumage, dominance and fluctuating asymmetry in female starlings. *Proc R Soc Lond B* 260:219–223
- Tella JL, Lemus JA, Carrete M, Blanco G (2008) The PHA test reflects acquired T-cell mediated immunocompetence in birds. *PloS One* 3:e3295
- Török J, Hegyi G, Garamszegi LZ (2003) Depigmented wing patch size is a condition dependent indicator of viability in male collared flycatchers. *Behav Ecol* 14:382–388
- Velando A, Lessells CM, Marquez JC (2001) The function of female and male ornaments in the Inca tern: evidence for links between ornament expression and both adult condition and reproductive performance. *J Avian Biol* 32:311–318
- Vinkler M, Schnitzer J, Munclinger P, Albrecht T (2012) Phytohaemagglutinin skin-swelling test in scarlet rose finch males: low-quality birds respond more strongly. *Anim Behav* 83:17–23
- Watson NL, Simmons LW (2010) Mate choice in the dung beetle *Onthophagus sagittarius*: are female horns ornaments? *Behav Ecol* 21:424–430
- Wilson RS, Condon CH, David G, FitzGibbon S, Niehaus AC, Pratt K (2010) Females prefer athletes, males fear the disadvantaged: different signals used in female choice and male competition have varied consequences. *Proc R Soc Lond B* 277:1923–1928
- Zuk M, Thornhill R, Ligon JD, Johnson K (1990) Parasites and mate choice in red jungle fowl. *Am Zool* 30:235–244